





HD COMMUNITY BIOREPOSITORY – ANTIBODY DATASHEET

Antibody Information	
CHDI#	CHDI-90000504
CCR# (Coriell ID)	CH00540 CH00552
Antibody Name	anti NTRK2-77:94, F8965
Description	Rabbit polyclonal antibody specific to residue 77-94 of Trk B
Target Protein Name	 <p>New anti-peptide TrkB antibodies</p> <p>61,62,63,64 (Q2-Q3)</p> <p>65,66 (T7-T8)</p> <p>68 (T8D-T9E)</p> <p>53,54 (D9D-T9E)</p> <p>358 (D9D-T9E)</p> <p>362 (D9D-T9E)</p> <p>55,56 (D9D-T9E)</p> <p>50,51,52 (D9D-T9E)</p> <p>54 (T9)</p> <p>Tyrosine Kinase</p> <p>Garcia Neuron 2007</p>
Volume/Concentration per vial	CH00540 1.408 mg/mL 250 µL CH00552 0.940 mg/mL 250 µL
Host Species	Rabbit
Isotype	Polyclonal
Immunogen/Epitope Sequence	KRLEIINEDDVEAYVGLK
Format	Affinity purified
Preparation	Borate-Buffered Saline + 1% BSA (w/v)
Source	Open Biosystems, Inc.
Date Made	12/17/2009 and 3/25/10
Species Reactivity	Mouse; Rat (as predicted by 100% homology of immunogen); Human=2 amino acid difference
Date Aliquotes Frozen	June 2010
Storage Conditions	-80°C
Application Notes	
Specificity Notes	strong immune-reactivity to TrkB, but not TrkA or TrkC
Validated Applications and Dilution*	WB: 1µg/mL ICC-FL: 1µg/mL (weak signal, appears hazy in appearance); IHC-Fz: 1:200 (mostly granule neurons in cerebellum) IP/ELISA: no data available
Supporting Data	<p>Courtesy of M. Chao (NYU):</p> <p>Rat E18 primary cortical neurons DIV6, starved 6h in MEM/glucose + 25 ng/mL BDNF for 10 min; 7.5 µg/mL lysate/lane; all antibodies used at 1 µg/mL; band @ 100kDa reflects the truncated TrkB species</p> <p>C14 = Santa Cruz antibody; TrkB = Upstate Antibody</p>  <p>CHDI-90000504 anti NTK2-77:94; F8965 CH00552 C14 TrkB</p> <p>250 150 100</p>

Supporting Data continued	<p>Rat E18 primary cortical neurons were fixed in 4% PFA, 20% sucrose in PBS for 10 min at RT. The fixative was quenched and cells were permeabilized with 0.1% TX-100 in PBS for 3 min, blocked and incubated with TrkB antibodies @1 µg/mL and 2° antibodies @ 1:1000 in blocking solution. Cells were washed and mounted in Mowiol488. Images were taken on an EclipseE800 microscope equipped with a 20X objective (Nikon), FITC and TRITC filters (HQ96170M and HQ96171M, Chroma Technology Corp) and an Axioacam HR camera. using Axiovision software (Zeiss Vision GmBH).</p>	
	<p>Mouse Embryo brain section (E18) were dehydrated in sucrose/OCT and prepared as frozen sections</p>	

* Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 µL or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.

Our antibodies have not been fully characterized and we cannot guarantee that the antibodies will work in any application other than ELISA against the immunizing peptide. Whenever possible, we will make data including suggested working dilutions available and the information provided should be used as a guideline only. It is

recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. Species reactivity information on the datasheets is provided for reference only.

We would greatly appreciate your feedback on the antibody, whether positive or negative.

For research use only; not for use as a diagnostic.