


HD COMMUNITY BIOREPOSITORY – ANTIBODY DATASHEET

CHDI-90000505 CH00541	
CH00341	
CH00553	
anti NTRK2-77:94, F8966	
	due 77-94 of Trk B
	1 4
61,63,63,64 68 190 179 NGF 19-C1 S53,54 (200 299) Tyrosime Kinase	
CH00541 1.31 mg/mL 250 μL	
CH00553 0.76 mg/mL 250 μL	
Rabbit	
Polyclonal	
KRLEIINEDDVEAYVGLK	
	gy of immunogen); Human=2 amino acid
-80°C	
strong immune-reactivity to TrkB, but not	TrkA or TrkC
WB: 1μg/mL	
ICC-FL: $1\mu g/mL$ (slightly punctuate in cell b	ody, axons appear hazy, mildly bright);
IHC-Fz: no data available	
IP/ELISA: no data available	
Courtesy of M. Chao (NYU):	CHDI-90000505
Rat E18 primary cortical neurons	anti NTK2-77:94; F8966
DIV6, starved 6h in MEM/glucose + 25 ng/mL BDNF for 10 min; 7.5 µg/mL lysate/lane; all antibodies	CH00553 C14 TrkB
. •	250
	150
•	100
C14 = Santa Cruz antibody; TrkB = Upstate Antibody	
	-
	The second secon
	CH00541 1.31 mg/mL 250 μL CH00553 0.76 mg/mL 250 μL Rabbit Polyclonal KRLEIINEDDVEAYVGLK Affinity purified Borate-Buffered Saline + 1% BSA (w/v) Open Biosystems, Inc. 12/17/2009 and 3/25/10 Mouse; Rat (as predicted by 100% homolo difference June 2010 -80°C strong immune-reactivity to TrkB, but not WB: 1μg/mL ICC-FL: 1μg/mL (slightly punctuate in cell be IHC-Fz: no data available IP/ELISA: no data available IP/ELISA: no data available Courtesy of M. Chao (NYU): Rat E18 primary cortical neurons DIV6, starved 6h in MEM/glucose + 25 ng/mL BDNF for 10 min; 7.5 μg/mL lysate/lane; all antibodies used at 1 μg/mL; band @ 100kDa reflects the truncated TrkB species C14 = Santa Cruz antibody;

Supporting Data continued

Rat E18 primary cortical neurons were fixed in 4% PFA, 20% sucrose in PBS for 10 min at RT. The fixative was quenched and cells were permeabilized with 0.1% TX-100 in PBS for 3 min, blocked and incubated with TrkB antibodies @1 $\mu g/mL$ and 2° antibodies @ 1:1000 in blocking solution. Cells were washed and mounted in Mowiol488. Images were taken on an EclipseE800 microscope equipped with a 20X objective (Nikon), FITC and TRITC filters (HQ96170M and HQ96171M, Chroma Technology Corp) and an Axiocam HR camera. using Axiovision software (Zeiss Vision GmBH).



^{*} Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 μ L or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.

Our antibodies have not been fully characterized and we cannot guarantee that the antibodies will work in any application other than ELISA against the immunizing peptide. Whenever possible, we will make data including suggested working dilutions available and the information provided should be used as a guideline only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. Species reactivity information on the datasheets is provided for reference only.

We would greatly appreciate your feedback on the antibody, whether positive or negative.

For research use only; not for use as a diagnostic.